



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

STATISTICAL REPORTS OF STATES AND CITIES OF THE UNITED STATES,
YEARLY AND MONTHLY.

CALIFORNIA.—Reports from 28 localities having an aggregate population of 857,150 show 1,153 deaths, including diphtheria 14, enteric fever 9, scarlet fever 1, whooping cough 4, and 166 from tuberculosis.

San Diego.—Month of January, 1905: Estimated population, 25,000. Total number of deaths, 46, including diphtheria, 1, and 13 from phthisis pulmonalis.

CONNECTICUT—*Stamford*.—Month of February, 1905: Estimated population, 18,000. Number of deaths not reported. No deaths from contagious diseases reported.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended February 18, 1905, from 67 observers, indicate that smallpox and cholera morbus were more than usually prevalent, and pneumonia, pleuritis, phthisis pulmonalis, inflammation of bowels, scarlet fever, enteric fever, erysipelas, remittent fever, intermittent fever, whooping cough, measles, puerperal fever, inflammation of brain, dysentery, meningitis, and cholera infantum were less than usually prevalent.

Meningitis was reported present at 3, whooping cough at 6, diphtheria at 31, measles at 35, scarlet fever at 53, enteric fever at 59, smallpox at 74, pneumonia at 79, and phthisis pulmonalis at 188 places.

MINNESOTA—*Duluth*.—Month of January, 1905. Estimated population, 70,000. Total number of deaths, 55, including 4 from tuberculosis.

TENNESSEE—*Nashville*.—Month of January, 1905. Estimated population, 83,520—white, 52,420; colored, 31,100. Total number of deaths, 197—white, 95; colored, 102—including diphtheria 3, enteric fever 3, whooping cough 4, and 27 from tuberculosis.

VIRGINIA—*Richmond*.—Month of January, 1905. Estimated population, 100,000—white, 62,250; colored, 37,750. Total number of deaths, 125—white, 70; colored, 55—including enteric fever 2, scarlet fever 1, and 26 from tuberculosis.